

Original research

Fish species composition and length-weight relationships in Onaç Creek (Burdur-Turkey)

Deniz INNAL^{1,*}, Duygu Ceren ÇAĞLAN¹, Filiz ÖZDEMİR²

¹Department of Biology, Mehmet Akif Ersoy University, 15100, Burdur, Turkey

²Department of Biology, Hacettepe University, 06800 Ankara, Turkey

*Corresponding author email: innald@yahoo.com

Abstract: This study describes the fish species composition and their length-weight relationships from Onaç Creek (Burdur-Turkey). A total of 498 individuals representing five fish species belongs to three families were caught by electrofishing techniques from May 2013 to June 2015. *Cyprinidae* was the most abundant family in terms of species number. *Cyprinidae* species comprised 350 (70.28%) of 498 specimens. Non indigenous fish species represents the highest abundance (98%). Topmouth gudgeon, *Pseudorasbora parva* was the most frequently captured species (43.57%), followed by eastern mosquitofish, *Gambusia holbrooki* (28.92%) and Prussian carp, *Carassius gibelio* (25.5%). The length-weight relationships for Prussian carp, Topmouth gudgeon and Eastern mosquitofish populations were estimated as $W = 0.0097 TL^{3.187}$, $W = 0.0061 TL^{3.26}$ and $W = 0.0083 TL^{2.995}$, respectively.

Keywords: Non indigenous, invasive, dominancy, threats, endemic

Citing: İnnal, D., Çağlan, D.C., & Özdemir, F., 2019. Fish Species Composition and Length-Weight Relationships in Onaç Creek (Burdur-Turkey). *Acta Biologica Turcica*, 32(3): 135-142.

Introduction

Anatolia is home to a large number of endemic fish species (Geldiay and Balık, 2007; Çiçek et al., 2015). Many of the endemic fishes in Anatolia are now directly under several threats, which are mostly induced by human activities (Fricke et al., 2007). Anthropogenic movements of invasive species are considered one of the main threats to aquatic biodiversity. Many papers have reviewed status and distribution of alien species in the inland waters of Turkey (İnnal and Erk'akan, 2006; Çetinkaya, 2006; Balık and Ustaoglu, 2006).

The Lakes District of Turkey (located at southern Mediterranean region) is characterized by great diversity of natural ecosystems. Compared to the other parts of Turkey, this region has relatively richer endemic aquatic fauna because of its habitat richness coming from geographical process (Balık, 1995; Demirsoy, 2002).

The presence of alien fish in Lakes district of Turkey has been recorded by Yeğen et al. (2006), Küçük et al. (2006), Özkök et al. (2007), Çınar et al. (2007), Küçük et al. (2012), İlhan et al. (2014), Küçük et al. (2016), Güçlü et al. (2017), Apaydın-Yağcı et al. (2018). Approximately ten species of exotic or translocated fish species have been introduced into Lakes District of Turkey. The following species have produced self-sustained populations and widespread: *Carassius gibelio* (Bloch, 1782), *Pseudorasbora parva* (Temminck and Schlegel, 1846), *Gambusia holbrooki* Girard, 1859. Biological properties are lacking for Onaç Creek fishes. This study concerns an investigation of the population structures and length-weight relationships of *C. gibelio*, *P. parva* and *G. holbrooki* living in Onaç Creek, Burdur, Turkey.

Material and Methods

Fish specimens were caught monthly interval with electrofishing equipment from three different locations of

Onaç Creek (Burdur) between May 2013 and June 2015. The Onaç Creek flows from the northern part of the Kestel Mountain, 1300 meters high and fed by small springs and melted snow water. It occupies Bucak plain and pour into the temporary Kestel Lake. The decline of the rains led to reduce the water of Onaç Creek and it completely dries in some years.

Fish specimens were collected and identified species level according to Freyhof and Özuluğ (2006) and Kottelat and Freyhof (2007). The abundance of species was according to Şişli (1996). Abundance % = $[\text{Ni}/\text{Nt}] \times 100$, (Ni: The number of specimens of the species; Nt: The total number of specimens). Fish samples were immediately transported to the laboratory. Size (total length cm TL; precision 1 mm) and weight (total weight W to nearest 0.1 g) were calculated. Length-weight relationship was calculated using the equation $W = aL^b$ (Pauly, 1984), where *a* is a coefficient related to body form and *b* is an exponent indicating isometric growth when equal to 3. The significance of the *b* value for each species was tested by t-test. The degree of association between the variables was computed by the determination coefficient, r^2 .

Results

A total of 498 individuals representing five natives and non native fish species was caught throughout the study. Fish specimens belonging to three families (Poeciliidae, Cyprinodontidae and Cyprinidae). *P. parva* was the most frequently captured species, followed by *G. holbrooki* and *C. gibelio*. Two native species (*Aphanius* sp. and *Pseudophoxinus ninae*) were sampled in small percentages (2 %).

Fish assemblage composition (species numbers, abundance and status) in Onaç Creek is given in Table 1. Length and weight frequency distributions and length-

weight relationships are shown in Figure 1. Total length and weight for specimens of *P. parva* ranged from 2.8 to 6.7 cm and 0.143 to 2.9 g, respectively. Among all individuals, 53.92% were in the size range of 4.0-5.2 cm, 23.50% were smaller than 4 cm, and 22.58% were longer than 5.2 cm. Length-weight relationships were calculated as $W = 0.0061L^{3.26}$ ($r^2=0.92$) by using the data of 217 *P. parva* specimens.

The specimens of *G. holbrooki* ranged from 0.7 to 4.9 cm in total length and 0.003 to 1.253 g in total weight. The total length of 86.80% of the total sampled specimens were range of 1.4-3.5 cm, 3.47% were smaller than 1.4 cm, and 9.72% were longer than 3.5cm. Length-weight relationships were estimated as $W = 0.0083L^{2.9956}$ ($r^2=0.96$) by using 144 specimens.

Total length and weight of the sampled specimens of *C. gibelio* ranged from 3.4 to 27.9 cm and from 0.5 to 470.4 g, respectively. Among all sampled specimens, 76.38% were in the size range of 11-23 cm, 3.94% were smaller than 11 cm, and 19.69% were longer than 23 cm. Length-weight relationships were found to be $W = 0.0097L^{3.1874}$ ($r^2=0.99$) from 127 specimens.

Discussion

Fishes in the Onaç Creek have been categorized as native and alien species. The alien species, with three species, represented the highest abundance (98 %). High abundance of alien species in Onaç Creek demonstrates antropogenical intervention. *Aphanius* sp. and *P. ninae* are endemic to Anatolian fish fauna and both have an extremely limited distribution. The Onaç Creek is also type locality of *P. ninae* (Freyhof and Özuluğ, 2006). It has been assessed as Critically Endangered (Freyhof, 2014). They are under considerable pressure from a range of anthropogenic activities as other native species.

Table 1. Fish assemblage composition (species, density and status)

Species	Common name	Family	Status	N	N%
<i>Pseudorasbora parva</i>	Topmouth gudgeon	Cyprinidae	Alien	217	43.6
<i>Gambusia holbrooki</i>	Eastern mosquitofish	Poeciliidae	Alien	144	28.9
<i>Carassius gibelio</i>	Prussian Carp	Cyprinidae	Alien	127	25.5
<i>Pseudophoxinus ninae</i>	Onaç Spring minnow	Cyprinidae	Endemic	6	1.2
<i>Aphanius</i> sp.	Killifish	Cyprinodontidae	Endemic	4	0.8
Total				498	100




Figure 1. Length and weight frequency distributions and length-weight relationships for *Carassius gibelio*, *Pseudorasbora parva* and *Gambusia holbrooki*

Table 2. Summary of estimated length weight parameters for *Carassius gibelio*, *Pseudorasbora parva* and *Gambusia holbrooki*. (TL: Total Length SL: Standart Length FL: Fork Length)

n	L type	L(cm) min-max	W (g) min-max	b	Locality	Author
<i>Gambusia holbrooki</i>						
224	TL	1.0-4.0	0.01-0.70	2.56	Seyhan Dam Lake	Alagöz Ergüden and Ergüden (2008)
215	TL	1.0- 3.9	*	2.61	Seyhan Dam Lake	Alagöz Ergüden and Gökso (2009)
58	SL	1.4-3.5	*	2.68	Sapanca Lake	Tarkan et al. (2009)
1582	TL	1.0-5.7	*	2.927	Seyhan Dam Lake	Alagöz Ergüden (2012)
5	TL	2.60- 3.9	0.20-0.80	2.945	Marmara Lake	İlhan and Sarı (2015)
909	TL	1.64-4.31	*	2.952	Lake Pamvotis	Gkenas et al.(2012)
25	TL	1.56-3.065	*	2.986	Sirzar River	Eagderi and Radkhah (2015)
144	TL	0.7-4.9	0.003-1.253	2.996	Onaç Creek	Present Study
35	TL	2.38-4.05	*	3.048	Iran	Esmaili and Ebrahimi (2006)
50	TL	2.25-3.66	*	3.086	Gamasiab River	Eagderi and Radkhah (2015)
4	TL	2.3-2.9	*	3.158	Porto Logos	Koutrakis and Tsikliras (2003)
162	?	1.15-5.8	0.01-2.62	3.168	Murray-Darling River System	Llewellyn (2011)
705	TL	1.3-5.5	0.02-2.31	3.17	Fethiye-Akgöl	Öztürk and İkiz (2004)
639	TL	1.3-5.8	0.02-5.83	3.27	Ortaca	Öztürk and İkiz (2004)
27	TL	1.4-4.41	*	3.31	Neretva Estuary	Dulcic and Glamuzina (2006)
60	FL	1.9 -4.9	*	3.37	Segura River tributaries	Andreu Soler et al. (2006)
682	TL	1.7-5.5	0.06-2.58	3.37	Dalaman	Öztürk and İkiz (2004)
671	TL	1.0-5.1	*	3.382	Rihios Estuary	Koutrakis and Tsikliras (2003)
15	TL	3.2-4.7	*	3.42	Büyükçekmece Dam Lake	Tarkan et al. (2006)
19	TL	2.0-4.4	*	3.49	Ömerli Dam Lake	Tarkan et al. (2006)
57	FL	2.0 -5.7	*	3.59	Segura River	Andreu Soler et al. (2006)
43	TL	2.57-5.53	*	3.763	Pond in Kashmar	Eagderi and Radkhah (2015)
119	FL	2.2 -5.7	*	3.81	Segura River reservoirs	Andreu Soler et al. (2006)

Table 2. continue

<i>Pseudorasbora parva</i>						
76	TL	4.9-12.0	*	2.9	Hunan H. (China)	Xie et al. (2015)
116	TL	5.20-11.0	1.60-14.60	2.929	Marmara Lake	İlhan and Sari (2015)
20	TL	5.46-8.37	1.7-6.6	3	İli River (China)	Sui et al. (2015)
33	TL	4.58-7.50	*	3.01	Iran	Esmaeili and Ebrahimi (2006)
141	TL	3.5-9.7	*	3.02	Tarim River (China)	Huo et al. (2012)
1329	FL	2.5-11.8	*	3.032	UK	Britton and Davies (2007)
36	TL	5.0-9.8	*	3.06	Saemangeum Reservoir (S.Korea)	Kim et al. (2015)
162	SL	2.9-9.5	*	3.08	Chishui	Liu et al. (2014)
373	TL	2.31-9.91	0.07-10.0	3.09	Lhasa R.B. (China)	Fan et al. (2015)
2674	TL	1.7-11.9	*	3.09	Erhai Lake (China)	Tang et al. (2013)
8	TL	4.0-6.6	*	3.091	Ergis River (China)	Huo et al. (2011)
30	SL	3.5-7.9	*	3.12	Tian –e-zhou	Wang et al. (2012)
25	TL	3.1-7.5	0.39-4.92	3.15	Zarrineh (Iran)	Radkhah and Eagderi (2015)
7815	TL	1.9-12.5	*	3.204	Flanders (Belgium)	Verreycken et al. (2011)
217	TL	2.8-6.7	0.143-2.9	3.26	Onaç Creek	Present Study
3368	TL	1.8-9.6	*	3.32	Hirfanlı Reservoir	Kirankaya et al. (2014)
30	TL	3.0-7.2	*	3.37	Sirwan River (Iran)	Hasankhani et al. (2014)
435	TL	5.1-12.0	*	3.55	Strymon River (Greece)	Petriki et al. (2011)
<i>Carassius gibelio</i>						
317	TL	10.7-31.0	26-450	2.571	Seyhan River	Alagöz Ergüden (2015a)
160	TL	11-29.30	40.10-412.9	2.65	Seyhan Dam Lake	Alagöz Ergüden (2015b)
344	FL	5.6-26.8	3.8 -597	2.80(♀♀) 3.05(♂♂)	Gelingüllü Dam Lake	Kirankaya and Ekmekçi (2013)
600	TL	21.9-37.0	*	2.81	Chimaditis Lake	Leonardos et al. (2008)
144	TL	12.2-39.5	26.31-445.5	2.856	Kızıllırmak River Basin	Sungur Birecikligil et al. (2016)
314	TL	13.0-35.0	50-900	2.866	Danube River	Gheorghe et al.(2012)
2325	FL	9.7-25.5	23.6-269	2.87	Buldan Dam Lake	Sarı et al. (2008)
95	TL	11.3-35.5	*	2.88	Anzali Wetland	Moradinassab et al. (2012)
480	TL	23.0–31.4	150.88- 622.02	2.886(♀♀) 2.981(♂♂)	İkizcetepeler Dam Lake	Erdoğan et al. (2014)
149	FL	14.8-32.5	43.1-807.3	2.94	Seyitler Reservoir	Bulut et al. (2013)
143	TL	12.9-32.3	*	2.945	Volvi	Bobori (2010)
2213	TL	6.80-27.50	4.90-372.20	2.974	Marmara Lake	İlhan and Sari (2015)
173	FL	16.9-30.0	125-730	2.978	Bafra Fish Lake	Bostancı et al. (2007)
93	TL	24.6-37.5	281.9-870	3.04	Ain Zada Reservoir	Mimeche and Biche (2015)
205	TL	8.4-30.7	*	3.059	Doirani	Bobori (2010)
102	TL	8.2-25.2	*	3.11	Volvi Lake	Kleandhitis et al. (1999)
128	TL	10.3-30.5	25-607	3.114	Aksu River	İnnal (2012)
3114	FL	7.8-32.2	5-829	3.125	İzник Lake	Uysal et al. (2015)
1717	FL	7.5-33.3	8.0-1073	3.128	Eğirdir Lake	Özkök et al. (2007)
487	FL	4.4-31.4	1.44-774.4	3.134	Büyükçekmece Dam Lake	Saç and Okgerman (2015)
155	FL	13.4-26.5	58-550	3.149	Ladik Dam Lake	Yazıcıoğlu et al. (2013)
616	FL	9.0-33.0		3.152	Eğirdir Lake	Balık et al. (2004)
730	TL	3-35.7	*	3.18	Ömerli Dam Lake	Tarkan et al. (2006)
*	TL	1.8-46.6	*	3.183	Flemish inland waters	Verreycken et al. (2011)
482	FL	7.1-27.4	6.0-495	3.186	Beyşehir Lake	Çınar et al. (2007)
49	TL	8.3-33.5	*	3.187	Mikri Prespa	Bobori (2010)
127	TL	3.4-27.9	0.5-470.4	3.187	Onaç Creek	Present Study
363	TL	5.2-30.2	*	3.25	İzник Lake	Tarkan et al. (2006)
15	TL	37.5-42.3	*	4.3001	Vransko Lake	Treer et al. (2011)

One of the main threats to its environment (Onaç Creek) is the introduction of the non native fish species [*P. parva*, *C. gibelio* and *G. holbrooki*] all of them are invasive. The ecology of native aquatic systems in Lakes

District of Turkey has changed considerably in the last decade due to human activities, which have resulted in decreased water quality and quantity (pers. comm. with local authorities). Date and agent of first introduction of

these alien fish species in Onaç Creek are unknown. *C. gibelio*, *P. parva* and *G. holbrooki* are continuing to steadily increase in abundance, and distribution in the lakes district of Turkey (İnnal D., pers. Observation; Yeğen et al., 2006; Küçük et al., 2009). The extensive geographic distribution and the range of habitats occupied by these invasive species suggest that their wide physiological tolerance. Strong competitive impacts of non-indigenous fishes on native species have been shown in freshwater systems (Özuluğ et al. 2005; Şaşı and Balık, 2003). Subsequent decline of many indigenous fishes was reported from one of the important lake system (Lake Eğirdir) of this region (Küçük et al., 2009).

Another potential risk factor is drought out which alter the normal river flow and may restrict the movement of the fish along the natural creek channel. In some periods Algae explosions have been observed. During these periods some *Aphanius sp.* and *P. ninae* individuals died. These deaths are thought to be caused by toxins from algae.

The length-weight relationships of three fish species belonging to two families were estimated. Comparison of length-weight relationships parameters of species between the present study and previous studies is shown in Table 2. In this study, the b value of the three species remained within the expected range of 2.5-3.5 (Froese, 2006). b value of *P. parva* (95 % confidence intervals 3.2473-3.2806; $p < 0.001$) and *C. gibelio* (95 % confidence intervals 3.1824-3.1953; $p < 0.001$) were significantly different from 3 and this indicate showed positively allometric growth, whereas *G. holbrooki* (95 % confidence intervals 2.9269-3.1285; $p > 0.05$) exhibited isometric growth. The slope b value of the LWR equations showed great variations from one population to another within the same species (Table 2). These variations in length-weight relationships could be caused by sampling and preservation protocols, environmental factors, food availability, diets, genetic factors, gonad maturities, sexes, parasitisms, seasons and other interspecific factors (Tesch, 1971; Wootton, 1998).

In conclusion, this study provides some basic information about three invasive fish species and is thought to be useful for future studies. The role of invasive fish species in the inland water systems of Turkey and their effects on local populations need to be the subject of future research.

Acknowledgements

We would like to thank Sevilay Karataş for her valuable contributions in laboratory studies.

References

- Alagöz-Ergüden S., Ergüden D. 2008. A preliminary research on growth feature of the mosquitofish *Gambusia affinis* (Baird and Girard, 1853) determined in Seyhan Dam Lake (Adana-Turkey). *Journal of Fisheries Sciences*, 2 (3): 312-320.
- Alagöz-Ergüden S., Göksu M.Z.L. 2009. Length-weight relationships for 12 fish species caught in Seyhan Dam Lake in southern Anatolia, Adana, Turkey. *Journal of Applied Ichthyology*, 25: 501-502.
- Alagöz-Ergüden S. 2012. Age, growth, sex ratio and diet of eastern mosquitofish *Gambusia holbrooki* Girard, 1859 in Seyhan Dam Lake. *Iran Journal of Fisheries Science*, 12: 204-218.
- Alagöz-Ergüden S. 2015a. Age and growth properties of Prussian Carp, *Carassius gibelio* Bloch, 1782) living in the middle basin of Seyhan River in Adana, Turkey. *Pakistan Journal of Zoology*, 47(5): 1365-1371.
- Alagöz-Ergüden S. 2015b. Determination of condition factor and length-weight relationship of the Prucian Carp, *Carassius gibelio* (Bloch, 1782) inhabiting Seyhan Dam Lake. *International Journal of Scientific and Technological Research*, 1(1): 157-166.
- Andreu-Soler A., Oliva-Paterna F.J., Torralva M. 2006. A review of length-weight relationships of fish from the Segura River basin (SE Iberian Peninsula). *Journal of Applied Ichthyology*, 022: 295-296.
- Apaydın-Yağcı M., Alp A., Yağcı, A., Koçer M.A.T. 2018. In book: Selected Studies in Biodiversity, Chapter 17: Prey Selection of *Pseudorasbora parva* (Temminck and Schlegel, 1846) in a Freshwater Ecosystem (Lake Eğirdir/Turkey). DOI: 10.5772/intechopen.70471
- Balık S. 1995. Freshwater fish in Anatolia, Turkey. *Biological Conservation*, 72: 213-223.
- Balık S., Ustaoglu M.R. 2006. Fish introducing studies in lakes, ponds and reservoirs of Turkey and their results. I. Symposium on Management of Reservoirs and Fish Stocking, MARA, Mediterranean Fisheries, Production and Education Institute, pp: 1-10.
- Balık I., Özkök R., Çubuk H., Uysal R. 2004. Investigation of some biological characteristics of the silver crucian carp, *Carassius gibelio* (Bloch 1782) population in Lake Eğirdir. *Turkish Journal of Zoology*, 28 (1):19-28.
- Bobori D.C., Moutopoulos D.K., Bekri M., Salvarina I., Munoz A.I.P. 2010. Length-weight relationships of freshwater fish species caught in three Greek lakes. *Journal of Biological Research-Thessaloniki*, 14: 219-224.

- Bostancı D., Polat N., Kandemir S., Yılmaz S. 2007. Length-weight relationship and condition factor of *Carassius gibelio* (Bloch, 1782) in Bafra Balık Lake. Süleyman Demirel University, Faculty of Art and Science, Journal of Science, 2: 117-125.
- Britton J. R., Davies G.D. 2007. Length-weight relationships of the invasive topmouth gudgeon (*Pseudorasbora parva*) in ten lakes in the UK. Journal of Applied Ichthyology, 23: 624-626.
- Bulut S., Mert R., Algan B., Özbek M., Ünal B., Konuk M. 2013. Several growth characteristics of an invasive cyprinid fish (*Carassius gibelio* Bloch, 1782). Notulae Scientia Biologicae, 5 (2): 133-138.
- Çetinkaya O. 2006. Exotic and native fish species that introduced or stocked into Turkish waters, their impacts on aquaculture, fisheries, wild populations and aquatic ecosystems: A preliminary study on constructing a database. Symposium on Management of Reservoirs and Fish Stocking, MARA, Mediterranean Fisheries, Production and Education Institute, pp: 205-235.
- Çınar S., Çubuk H., Özkök R., Tümgelir L., Çetinkaya S., Erol K.G., Ceylan M. 2007. Beyşehir Gölü'ndeki gümüşü havuz balığı (*Carassius gibelio* Bloch, 1782) populasyonunun büyüme özellikleri. Turkish Journal of Aquatic Life, 3-5 (5-8): 401-409.
- Çiçek E., Sungur Birecikligil S., Fricke R. 2015. Freshwater fishes of Turkey: A revised and updated annotated checklist. Biharean Biologist, 9 (2): 141-157.
- Demirsoy A. 2002. General Zoogeography and Turkish Zoogeography. Meteksan publishing, Ankara.
- Dulcic J., Glamuzina B. 2006. Length-weight relationships for selected fish species from three eastern Adriatic estuarine systems (Croatia). Journal of Applied Ichthyology, 22: 254-256.
- Eagderi S., Radkhah A. 2015. Length-weight relationship and condition factor of mosquitofish (*Gambusia holbrooki*) in three inland basins of Iran. Poeciliid Research, 5 (1): 39-43.
- Erdoğan Z., Torcu-Koç H., Güngör S., Ulunehir G. 2014. Age, growth and reproductive properties of an invasive species *Carassius gibelio* (Bloch, 1782) (Cyprinidae) in the İkizcetepeler Dam Lake (Balıkesir), Turkey. Periodicum Biologorum, 116 (3): 285-291.
- Esmaceli H.R., Ebrahimi M. 2006. Length-weight relationships of some freshwater fishes of Iran. Journal of Applied Ichthyology, 22(4): 328-329.
- Fan L.Q., Zhang X.J., Pan G. 2015. Length-weight and length-length relationships for nine fish species from Lhasa River Basin, Tibet, China. Journal of Applied Ichthyology, 31: 807-808.
- Freyhof J., Özüluğ M. 2006. *Pseudophoxinus ninae*, a new species from Central Anatolia, Turkey (Teleostei: Cyprinidae). Ichthyological Exploration of Freshwaters. 17 (3): 255-259.
- Freyhof, J. 2014. *Pseudophoxinus ninae*. The IUCN Red List of Threatened Species 2014: e.T19449277A19849704. Available from: <http://dx.doi.org/10.2305/IUCN.UK.20141.RLTS.T19449277A19849704.en>, Retrieved 18/06/ 2018.
- Fricke R., Bilecenoğlu M., Sarı, H.M. 2007. Annotated checklist of fish and lamprey species (Gnathostoma and Petromyzontomorphi) of Turkey, including a Red List of threatened and declining species. Stuttgarter Beiträge zur Naturkunde A (Biologie), 706: 1-172.
- Froese R. 2006. Cube law, condition factor and weight-length relationships: history, meta-analysis and recommendations. Journal of Applied Ichthyology, 22: 241-253.
- Geldiay R., Balık S. 2007. Freshwater Fishes of Turkey. Ege University Faculty of Fisheries Publishing, No:46, İzmir, Turkey.
- Gheorghe D.C., Nica A., Cristea V., Răzlog G.P. 2012. Growth and mortality estimation parameters for the Prusian carp (*Carassius gibelio*, Bloch, 1782) population from Danube River (km 170-196). UASVM Iasi, Lucrări Ştiinţifice Journal, Seria Zootehnie, 57 (17): 164-169.
- Gkenas C., Oikonomou A., Economou A., Kiosse F., Leonardos I. 2012. Life history pattern and feeding habits of the invasive mosquitofish, *Gambusia holbrooki*, in Lake Pamvotis (NW Greece). Journal of Biological Research, 17: 121-136.
- Güçlü S.S., Erdoğan Ö. 2017. Age, growth, sex ratio and feeding of *Knipowitschia caucasica* (Berg, 1916) (Actinopterygii, Gobiidae) in non-native species of Eğirdir Lake (Turkey). Acta Biologica Turcica, 30 (1): 1-6.
- Hasankhani M., Keivany Y., Daliri M., Pouladi M., Soofiani N.M. 2014. Length-weight and length-length relationships of four species (*Barbus lacerta* Heckel, 1843), *Oxynoemacheilus angorae* (Steindachner, 1897), *Squalius lepidus* (Heckel, 1843) and *Pseudorasbora parva* (Temminck & Schlegel, 1846) from the Sirwan River (western Iran). Journal of Applied Ichthyology, 30 (1): 206-207.
- Huo T.B., Yuan M.Y., Jiang, Z. 2011. Length-weight relationships of 23 fish species from the Ergis River in Xingjiang, China. Journal of Applied Ichthyology, 27 (3): 937-938.
- Huo T.B., Jiang Z.F., Karjan A., Wang Z.C, Tang F.J., Yuo H.X. 2012. Length-weight relationships of 16 fish species from the Tarim River, China. Journal of Applied Ichthyology, 28: 152-153.
- İlhan A., Balık S., Sarı H.M. 2014. Recent distribution and conservation status of endemic freshwater fishes from central and western Anatolia. Istanbul University Journal of Fisheries & Aquatic Sciences, 29 (2): 9-34.

- İlhan A., Sarı H. M. 2015. Length-weight relationships of fish species in Marmara Lake, West Anatolia, Turkey. *Croatian Journal of Fisheries*, 73 (1): 30-32.
- İnnal D., Erk'akan F. 2006. Effects of exotic and translocated fish species in the inland waters of Turkey. *Reviews in Fish Biology and Fisheries*, 16: 39-50.
- İnnal D. 2012. Age and growth properties of *Carassius gibelio* (Cyprinidae) living in Aksu River Estuary (Antalya-Turkey). *Review of Hydrobiology*, 5 (2):97-109.
- Kırankaya Ş.G., Ekmekçi F.G. 2013. Life-history traits of the invasive population of Prussian Carp, *Carassius gibelio* (Actinopteri: Cypriniformes: Cyprinidae), from Gelingüllü Reservoir, Yozgat, Turkey. *Acta Ichthyologica Et Piscatoria*, 43 (1): 31-40.
- Kırankaya S.G., Ekmekçi F.G., Yalçın-Özdilek S., Yoğurtcuoğlu B., Gençoğlu, L. 2014. Condition, length-weight and length-length relationships for five species from Hirfanlı reservoir, Turkey. *Journal of Fisheries Sciences*, 8: 208-213.
- Kim, J.H., Yoon J.D., Won D.H., Byeon M.S., Jang M.H. 2015. Length-weight relationships of 19 fish species from Saemangeum reservoir in South Korea. *Journal of Applied Ichthyology*, 31: 951-953.
- Kleantidis P.K., Sinis A.I., Stergiou K.I. 1999. Length-weight relationships of freshwater fishes in Greece. *Naga, ICLARM Quarterly*, 22 (4): 37-41.
- Kottelat M., Freyhof J. 2007. *Handbook of European freshwater fishes*. Publications Kottelat, Cornol and Freyhof, Berlin. ISBN: 978-2-8399-0298-4., 646 p.
- Koutrakis, E.T., Tsikliras A.C. 2003. Length-weight relationships of fishes from three northern Aegean estuarine systems (Greece). *Journal of Applied Ichthyology*, 19: 258-260.
- Küçük F., Gülle İ., Güçlü S.S., Gümüş E., Demir O. 2006. Eğirdir Gölü'ne Sonradan Giren Gümüşbalığı (*Atherina boyeri* RISSO, 1810)'nın Göl Ekosistemine ve Balıkçılığa Etkisi, 1. Balıklandırma ve Rezervuar Yönetimi Sempozyumu, Bildiriler Kitabı, 119-128 s., 7-9 Şubat 2006, Antalya.
- Küçük F., Sarı H.M., Demir O., Gülle İ. 2009. Review of the ichthyofaunal changes in Lake Eğirdir between 1915 and 2007. *Turkish Journal of Zoology*, 33: 277-286.
- Küçük F., Güçlü S.S., Gülle İ., Güçlü Z., Çiçek N.L., Diken G. 2012. Reproductive Features of Big Scale-Sand Smelt, *Atherina boyeri* (Risso, 1810), an Exotic Fish in Lake Eğirdir (Isparta, Turkey), *Turkish Journal of Fisheries and Aquatic Sciences* 12: 1-2.
- Küçük F., Gülle İ., İnnal D., Güçlü S.S., 2016. Burdur ili balıkları: Biyoçeşitlilik envanteri, populasyon ve habitat durumları, Mehmet Akif Ersoy Üniversitesi Fen Bilimleri Enstitüsü Dergisi, 7: 232-239.
- Leonardos I.D., Tsikliras A.C., Eleftheriou V., Cladas Y., Kagalou I., Chortatou R., Papigiotti O. 2008. Life history characteristics of an invasive cyprinid fish (*Carassius gibelio*) in Chimaditis Lake (northern Greece). *Journal of Applied Ichthyology*, 24: 213-217.
- Liu F., Cao W., Wang J. 2014. Length-weight relationships of 77 fish species from the Chishui River, China. *Journal of Applied Ichthyology*, 30: 254-256.
- Llewellyn L. 2011. Length-weight relationships of freshwater fish from the Murray-Darling River System in inland New South Wales, with particular reference to the golden perch *Macquaria ambigua*. *Australian Zoologist*, 35: 435-457.
- Mimeche F., Biche M. 2015. Length-weight relationships of four non-native cyprinid from the semiarid region in North-East of Algeria. *AAFL Bioflux*, 8 (1): 82-88.
- Moradinasab G.H., Daliri M., Ghorbani R., Paighambari S.Y., Davoodi R. 2012. Length-weight and length-length relationships, Relative condition factor and Fulton's condition factor of five cyprinid species in Anzali wetland, southwest of the Caspian Sea. *Caspian Journal of Environmental Sciences (CJES)*, 10: 25-31.
- Özkök R., Çubuk H., Tümgelir L., Uysal R., Çınar S., Küçükpara R., Erol G.K., Ceylan M. 2007. Growth features of silver crusian carp *Carassius gibelio* population in Lake Eğirdir. *Turkish Journal of Aquatic Life*, 3-5 (5-8): 313-322.
- Öztürk S., İkiz R. 2004. Some biological properties of mosquitofish populations (*Gambusia affinis*) living in inland waters of the western Mediterranean region of Turkey. *Turkish Journal of Veterinary and Animal Sciences*, 28 :355-361.
- Özuluğ M., Acıpınar H., Gaygusuz O., Gürsoy C., Tarkan A.S. 2005. Effects of human factor on the fish fauna in a drinking-water resource (Omerli Dam Lake-Istanbul, Turkey). *Research Journal of Agriculture and Biological Sciences*, 1 (1): 50-55.
- Pauly D. 1984. *Fish population dynamics in tropical waters: a manual for use with programmable calculators*. ICLARM Studies and Reviews 8. International Center for Living Aquatic Resources Management, Philippines, Manila, pp: 325.
- Petriki O., Gousia E., Bobori D.C. 2011. Weight-length relationships of 36 fish species from the River Strymon system (northern Greece). *Journal of Applied Ichthyology*, 27 (3): 939-941.
- Radkhah A., Eagderi S. 2015. Length-weight and length-length relationships and condition factor of six cyprinid fish species from Zarrineh River (Urmia Lake basin, Iran) Iran. *Journal of Ichthyology*, 2 (1): 61-64.
- Saç G., Okgerman H. 2015. Growth and reproduction of a non-native fish species *Carassius gibelio* (Bloch, 1782) from Büyükçekmece Lake (İstanbul, Turkey). *IUFS Journal of Biology*, 74 (1): 1-12.

- Sarı M.H., Balık S., Ustaoglu M.R., İlhan A. 2008. Population structure, growth and mortality of *Carassius gibelio* (Bloch, 1782) in Buldan Dam Lake. Turkish Journal of Fisheries and Aquatic Sciences, 8: 25-29.
- Şaşı H., Balık S. 2003. Age, growth and sex ratio of chub (*Leuciscus cephalus* L., 1758) in Topcam Dam Lake (Aydın). Ege University Journal of Fisheries and Aquatic Sciences, 20 (3-4): 503-515.
- Şişli M.N. 1996. Ekoloji, Çevre Bilim, Yeni Fersa Publishing, Ankara, pp. 492.
- Sui, X.Y., Li X.Q., Sun H.Y., Chen Y.F. 2015. Length-weight relationship of 13 species from the Ili River, China. Journal of Applied Ichthyology, 31: 1155-1157.
- Sungur Birecikligil S., Çiçek E., Öztürk S., Seçer B., Celepoğlu Y. 2016. Length-length, length-weight relationship and condition factor of fishes in Nevşehir Province, Kızılırmak River Basin (Turkey), Acta Biologica Turcica, 29 (3): 72-77.
- Tang J.F., Ye S.W., Liu J.S., Zhang T.L., Zhu F.Y., Guo Z.Q., Li Z.J. 2013. Composition and length-weight relationships of fish species in Lake Erhai, southwestern China. Journal of Applied Ichthyology, 29: 1179-1182.
- Tarkan A.S., Gaygusuz O., Acıpınar H., Gürsoy C., Özuluğ M. 2006. Length-weight relationship of fishes from the Marmara region (NW-Turkey). Journal of Applied Ichthyology, 22: 271-273.
- Tarkan A.S., Gaygusuz O., Özuluğ M., Gürsoy-Gaygusuz C., Saç G. 2009. Length-weight relationships of six freshwater fishes from the small creeks flowing into Lake Sapanca, NW Turkey. Journal of Applied Ichthyology, 25: 230-231.
- Tesch, W. 1971. Age and growth. In: W.E. Ricker (Ed.). Methods for assessment of fish production in fresh waters, 2nd edn. International Biological Programme, and Blackwell Scientific Publications. Oxford and Edinburg, UK. pp: 97-130.
- Treer T., Matulic D., Bogdanovic G., Anicic I., Safner R., Piria M. 2011. The condition of allochthonous fishes in the Mediterranean Vransko Lake. Journal of Applied Ichthyology, 27: 965-967.
- Uysal R., Alp A., Yeğen V., Apaydın-Yağcı M., Çetinkaya S., Yağcı A., Bostan H., Cesur M., Küçükpara R. 2015. The growth properties of Prussian Carp (*Carassius gibelio* Bloch, 1782) population in İznik Lake (Bursa/Turkey). Limnofish-Journal of Limnology and Freshwater Fisheries Research, 1 (1): 19-27. (in Turkish)
- Verreycken H.G., Thuyne V., Belpaire C. 2011. Length-weight relationships of 40 freshwater fish species from two decades of monitoring in Flanders Belgium. Journal of Applied Ichthyology, 27 (6): 1416-1421.
- Wang T, Wang H.S., Sun G.W., Huang D., Shen J.H. 2012. Length-weight and length-length relationships for some Yangtze River fishes in Tian-e-zhou Oxbow, China. Journal of Applied Ichthyology, 28: 660- 662.
- Wootton R.J. 1998. Ecology of teleost fishes. Kluwer Academic Publishers, Dordrecht, the Netherlands, 396 p.
- Xie J.Y., Kang Z.J., Yang J., Yang D.D. 2015. Length-weight relationships for 15 fish species from the Hunan Hupingshan National Nature Reserve in central China. Journal of Applied Ichthyology, 31: 221-222.
- Yazıcıoğlu O., Yılmaz S., Yazıcı R., Polat N. 2013. Condition factor, length-weight and length-length relationships of *Carassius gibelio* (Bloch, 1782) in Ladik Lake (Samsun, Turkey). The Black Sea Journal of Sciences, 3: 72-80. (in Turkish)
- Yeğen V., Balık S., Bostan H., Uysal R., Bilcen E. 2006. Göller bölgesindeki bazı göl ve baraj göllerinin balık faunalarının son durumu, 1. Ulusal Balıklandırma ve Rezervuar Yönetimi Sempozyumu, 7-9 Şubat 2006, Antalya. pp: 129-139.